



# Building Blocks of Clinical Practice

*Helping Athletic Trainers Build a Strong Foundation*



## Issue #2: Bacterial Infections of the Skin

### Folliculitis

#### Definition:

- Inflammation of a hair follicle
- Can progress down hair follicle or into multiple follicles and result in a furuncle or carbuncle

#### Causes:

- Bacterial or viral infection, chemical irritation
- Secondary to skin injury, which introduces bacteria to the area
- Shaving hairy areas may facilitate infection
- Occlusion of hair-bearing areas may facilitate growth of microbes
- Friction
- Hyperhidrosis

#### Symptoms:

- Areas are usually non-tender or slightly tender
- Area may itch
- Erythematous perifollicular papules or pustules may develop
- Grouped lesions

#### Types:

- Most common form is staphylococcal folliculitis

#### Diagnosis

- Confirmed by laboratory findings

#### Treatment:

- Antiseptic washes
- Warm compress
- Topical antibiotics
- Moderate and severe cases may require a systemic antibiotic

#### Prevention:

- Wash areas with antibacterial soap or benzoyl peroxide
- Practice good hygiene
- Avoid sharing razors and towels

### Carbuncles

#### Definition:

- A complication of folliculitis, a carbuncle is several furuncles that have merged
- Carbuncles are readily transmitted by skin-to-skin contact. Often, the direct cause of a carbuncle cannot be determined

#### Causes:

- Most carbuncles are caused by the bacteria staphylococcus aureus. The infection is contagious and may spread to other areas of the body or other people

#### Symptoms:

- A carbuncle is a swollen lump or mass under the skin which may be the size of a pea or as large as a golf ball
- The carbuncle may be red and irritated and might hurt when you touch it
- Pain gets worse as it fills with pus and dead tissue
- Other signs and symptoms include itching at the site of infection, skin inflammation around the wound, general ill feeling, fever, or fatigue. Pain improves as it drains

#### Diagnosis:

- Diagnosis is primarily based on the appearance of the skin
- Skin or mucosal biopsy culture may show staphylococcus or other bacteria

#### Treatment:

- Use of a warm, moist cloth several times a day on the carbuncle helps it to drain, which speeds healing
- Never squeeze a boil or attempt to cut it open because this can spread the infection and make the condition worse. It may drain on its own but more often than not a medical professional will drain it for the patient
- Once open, keep clean and use topical antibiotic ointments and keep covered



Picture reprinted with permission from [www.aoca.org](http://www.aoca.org).

© Jere Mammino, DO

*Continued...*

**NATA Research & Education Foundation**  
2952 Stemmons Freeway, Suite 200  
Dallas, TX 75247  
Phone: (214) 637-6282  
[www.natafoundation.org](http://www.natafoundation.org)

## Issue #2: Bacterial Infections of the Skin

*continued...*

### Impetigo

#### Definition:

- A bacterial infection caused by both streptococci and staphylococci species

#### Causes:

- Infects athletes with close skin to skin contact

#### Symptoms:

- Well defined, erythematous, yellow, crusted plaques distributed on the extremities and the head and neck

#### Diagnosis:

- Impetigo is usually diagnosed on the basis of clinical findings
- Bacterial culture and sensitivity are recommended (1) in cases to identify methicillin-resistant *Staphylococcus aureus* (MRSA), (2) if an outbreak of impetigo has occurred, or (3) if poststreptococcal glomerulonephritis is present
- Exudate from underneath the crust is sent for culture

#### Treatment:

- Both topical and oral antibiotics. Individuals should be kept from sporting activities that involve skin to skin contact until 5 days after the initiation of therapy



Picture reprinted with permission from [www.impetigodoctor.com](http://www.impetigodoctor.com)

### Furuncles/Boils

#### Definition:

- A bacterial infection (typically *Staphylococcus aureus*) of the hair follicle

#### Causes:

- Furunculosis and carbunculosis are often caused by bacteria (germs) called *Staphylococcus aureus* or Staph
- The Staph bacteria are commonly found on the skin, between the buttocks (rear end) and groin, and in the nose

#### Symptoms:

- Initially, a red, swollen nodule that is exquisitely tender
- The lesion eventually develops a yellow, pus-filled center that may have drainage. The patient may also develop a fever

#### Diagnosis:

- The diagnosis is made based on symptoms
- The drainage should be cultured to determine appropriate antibiotic therapy

#### Treatment:

- Initially, warm soaks will allow lesion to drain spontaneously
- Large or unresponsive lesions should be surgically lanced
- Systemic antibiotics may be prescribed for large lesions and for lesions on the face

#### References:

Adams, B.B. (2002). *Dermatological disorders of the Athlete*. Sports Medicine, 32(5), 309-321.

Anderson, Hall and Martin, *Foundations of Athletic Training*, 3rd edition, 2005. p 738.

Cuppett and Walsh, *General Medical Conditions in the Athlete*, 2005, p 328-29.

Ferri FF, Ellis J, Jones R. Folliculitis. MD Consult – First Consult. Retrieved December 4, 2009. [http://proxy2.atsu.edu:2053/das/pdxmd/body/174549765-4/926364596?type=med&eid=9-u1.0-\\_1\\_mt\\_1014864](http://proxy2.atsu.edu:2053/das/pdxmd/body/174549765-4/926364596?type=med&eid=9-u1.0-_1_mt_1014864)

Wolff K, Johnson RA. *Fitzpatrick's Color Atlas and Synopsis of Clinical Dermatology*, 6th Edition. New

York: McGraw-Hill Medical, 2009. Retrieved via StatRef, December 4, 2009.

Stulburg DL, Penrod MA, Blanty RA. Common Bacterial Skin Infections. *American Family Physician*. 2002;66(1):119-124.

